



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Neuropathic changes in equine laminitis pain

Citation for published version:

Jones, E, Vinuela-Fernandez, N, Eager, RA, Delaney, A, Anderson, H, Patel, A, Robertson, DC, Allchorne, A, Sirinathsinghji, EC, Milne, EM, Macintyre, N, Shaw, DJ, Waran, NK, Mayhew, J & Fleetwood-Walker, SM 2007, 'Neuropathic changes in equine laminitis pain', *Pain*, vol. 132, no. 3, pp. 321-31.
<https://doi.org/10.1016/j.pain.2007.08.035>

Digital Object Identifier (DOI):

[10.1016/j.pain.2007.08.035](https://doi.org/10.1016/j.pain.2007.08.035)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Pain

Publisher Rights Statement:

© 2007 International Association for the Study of Pain. Published by Elsevier Inc. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Neuropathic changes in equine laminitis pain

Emma Jones^b, Ignacio Vinuela-Fernandez^a, Rachel A. Eager^b, Ada Delaney^a, Heather Anderson^a, Anisha Patel^a, Darren C. Robertson^a, Andrew Allchorne^a, Eva C. Sirinathsinghji^a, Elspeth M. Milne^b, Neil MacIntyre^b, Darren J. Shaw^b, Natalie K. Waran^b, Joe Mayhew^b and Susan M. Fleetwood-Walker^{a,*}

EJ and IVF contributed equally to the work.

^aCentre for Neuroscience Research, Division of Veterinary Biomedical Sciences, R(D)SVS

^bDivision of Veterinary Clinical Sciences, R(D)SVS, The University of Edinburgh, Roslin, Midlothian EH25 9RG, United Kingdom.

Abbreviated title: Neuropathic pain induced by equine laminitis

Number of text pages: 29; number of Figures: 4; number of Tables: 2.

Address for correspondence:

* Professor SM Fleetwood-Walker,
Centre for Neuroscience Research,
Division of Veterinary Biomedical Sciences,
University of Edinburgh,
Summerhall,
Edinburgh EH9 1QH
Scotland, UK.

Tel.: +44 131-6506091

Email address: s.m.fleetwood-walker@ed.ac.uk

Abstract

Laminitis is a common debilitating disease in horses that involves painful disruption of the lamellar dermo-epidermal junction within the hoof. This condition is often refractory to conventional anti-inflammatory analgesia and results in unremitting pain, which in severe cases requires euthanasia. The mechanisms underlying pain in laminitis were investigated using quantification of behavioural pain indicators in conjunction with histological studies of peripheral nerves innervating the hoof. Laminitic horses displayed consistently altered or abnormal behaviours such as increased forelimb lifting and an increased proportion of time spent at the back of the box compared to normal horses. Electron micrographic analysis of the digital nerve of laminitic horses showed peripheral nerve morphology to be abnormal, as well as having reduced numbers of unmyelinated (43.2%) and myelinated fibers (34.6%) compared to normal horses. Sensory nerve cell bodies innervating the hoof, in cervical, C8 dorsal root ganglia (DRG), showed an upregulated expression of the neuronal injury marker, activating transcription factor-3 (ATF3) in both large NF-200-immunopositive neurons and small neurons that were either peripherin- or IB4-positive. A significantly increased expression of neuropeptide Y (NPY) was also observed in myelinated afferent neurons. These changes are similar to those reported in other neuropathic pain states and were not observed in the C4 DRG of laminitic horses, which is not associated with innervation of the forelimb. This study provides novel evidence for a neuropathic component to the chronic pain state associated with equine laminitis, indicating that anti-neuropathic analgesic treatment may well have a role in the management of this condition.

Keywords: Neuropathic pain, Dorsal root ganglion, Equine laminitis, neuronal injury marker, ATF3

1. Introduction

Laminitis is a common cause of equine lameness involving one or more feet (Dyson, 2003). It is characterised by disruption of the dermo-epidermal laminar bond within the hoof (Fig. 1b, c) and subsequent structural weakness that can result in displacement of the pedal bone within the hoof capsule (Pollitt et al., 1998). The pathogenesis of this disease is poorly understood but it is generally thought that vascular disturbances leading to ischemia-reperfusion injury of the lamellar structures are involved in the pathophysiology of laminitis (Hood, 1999). Currently, no therapeutic regime is able to arrest or prevent its onset (Pollitt, 2003). Moreover, laminitic pain can be difficult to control using traditional anti-inflammatory agents and euthanasia on welfare grounds is not uncommon (Herthel and Hood, 1999; Swanson, 1999; Pollitt, 2003). Therefore, improved understanding of laminitis is much needed.

We hypothesized that the pathological inflammatory processes affecting the hoof laminae during laminitis also damage the sensory neurons innervating this region. Peripheral nerve injury can be associated with the generation of a neuropathic pain state characterised by allodynia (the perception of normally innocuous stimuli as painful), hyperalgesia (a heightened response to painful stimuli), spontaneous pain and a lack of response to conventional analgesics. A key factor in the neural plasticity underlying neuropathic (compared to inflammatory) pain is altered gene expression in sensory DRG neurons (Hökfelt et al., 1994; Cummins et al., 2000; Woolf and Salter, 2000; Xiao et al., 2002). This can be demonstrated by an increase in expression of the neuronal injury marker ATF3, a member of the activating transcription factor/cAMP-responsive element binding protein (ATF/CREB) family, in sensory DRG cells (Hai et al., 1999; Tsujino et al., 2000). Furthermore, phenotypic changes occur in primary afferent DRG neurons after peripheral nerve damage, resulting in altered expression of neuropeptides, including neuropeptide Y (NPY), the expression of which is induced from normally low levels in large diameter, neurofilament-200 (NF-200)-positive neurons following axotomy (Wakisaka et al., 1991; Hokfelt et al., 1994), nerve injury

(Ma and Bisby, 1998; Munglani et al., 1995), demyelination (Wallace et al., 2003) and streptozotocin-induced diabetes (Rittenhouse et al., 1996).

Injury to sensory nerves induces neurochemical, physiological and anatomical modifications to afferent and central neurons that are likely to contribute to chronic, sensitised neuropathic pain responses (Woolf and Salter, 2000). Such changes to the sensory neurons innervating the equine foot could lead to a clinically relevant component of chronic pain as it would explain the limited effectiveness of conventional analgesics in the treatment of laminitic pain (Herthel and Hood, 1999).

Therefore, the aims of this study were to identify and quantify equine laminitic pain using objective behavioural assessment, characterise peripheral nerve damage in the lateral digital nerve and demonstrate potential nerve injury-associated alterations in protein expression in DRG sensory neurons innervating the feet of horses with laminitis.

2. Methods

2.1. Behavioural observations in laminitic and clinically normal horses

In order to define and quantify the behavioural characteristics of equine laminitis, we carried out continuous video monitoring over 3 days to compare behaviours in laminitic and normal horses.

Seven horses admitted for management of refractory laminitis were selected using the following clinical criteria: animals must have displayed clinical signs consistent with this disease including multi-limb lameness, increased amplitude of the digital pulses, warmth across the dorsal hoof wall and a laminitic gait (Stashak, 2002). For details of all laminitic horses used in this study see Table 1.

Latero-medial radiographs of the fore limb digits were obtained from each of these horses (Butler et al, 2000). The position of the pedal bone within the hoof capsule was evaluated both subjectively and objectively by an experienced equine clinician using standard measures (Fig.1a).

111

112 Informed client consent was obtained in writing prior to the onset of data collection.
113 Laminitic horses received phenylbutazone (PBZ) twice daily at 08:00h and 20:00h
114 (Equipalazone Arnolds, UK; 2.2-4.0 mg.kg⁻¹) and intramuscular acepromazine three times
115 daily at 08:00h, 16:00h and 24:00h (ACP Novartis, UK; 0.02-0.04 mg.kg⁻¹). On the day of
116 admission to hospital the timing of drug administration varied between individuals. Pedal
117 bone support (Styrofoam Solar Support System™/Lilypads™) was provided at the clinician's
118 discretion. Subjects participated in the study for a maximum of 3 days. Seven age, type and
119 sex-matched horses, which were considered 'pain free' (control group) were stabled directly
120 opposite the laminitic horses and recorded simultaneously in order to account for extraneous
121 effects on behaviour. All horses were maintained on shavings and had free access to water.
122 Laminitic animals were fed restricted rations of soaked hay, as is standard procedure, whereas
123 control animals received haylage *ad libitum*.
124 24-hour time-lapse video equipment (AG-6124, Panasonic) was used to record undisturbed
125 behaviour in each stable. Point samples of 1 hour duration were taken at 8 hour-intervals, at
126 06:00h, 14:00h and 22:00h, during 3 days starting at 14.00h on Day 1. Samples were analysed
127 continuously for duration of state and frequency of event behaviour (The Observer™ vs. 4.1,
128 Noldus Information Technology, The Netherlands).
129 Two behaviours were selected for statistical analysis as being representative of the behaviours
130 where changes were most likely to be observed (Price et al., 2003; Reitmann et al., 2004).
131 Frequency of 'forelimb lifting' (as lifts min⁻¹) was defined as the raising and lowering of a
132 forelimb, without locomotion and was adjusted for total time spent standing. 'Proportion of
133 time spent at the back of the box' was defined as time spent positioned in the furthest 50% of
134 the box, away from the entrance to the stable.
135 On a repeated dose regime, peak PBZ concentrations occur between 2 and 6 hours following
136 administration, although individual variation is high (Gerring et al., 1981). In the present
137 study a 12-hour dosing regime was used, minimising variation and increasing the probability
138 of the maintenance of a 'steady state'. For sample point analysis of behaviours, data were

collected at three different time points to reflect an expected minimum plasma PBZ concentration (06:00h) and shortest (2 hours post-administration – 22:00h) and longest (6 hours post administration – 14:00h) times for peak PBZ concentrations were chosen. In addition, to evaluate some of the possible effects of drug accumulation, analyses were repeated just using data from day 1, days 1 and 2 and days 1, 2 and 3.

2.2. Morphological investigations

Lateral digital nerves were obtained from five horses euthanised on clinical grounds due to laminitis which was either recurrent or refractory to therapy and also from four horses which had no history of forelimb lameness that were euthanised for clinical reasons other than forelimb pathology (control group). The lateral digital nerves were removed from the forelimb (3 cm long segments) at the level of the proximal sesamoid bone and fixed for 4 hours in 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium cacodylate buffer, pH 7.3, post-fixed in OsO₄, and embedded in Araldite. For light microscopy, 1 µm resin sections of the nerve were stained with Toluidine blue and three fascicles were chosen at random by bright-field microscopy. Ultra-thin (80 nm) sections were stained with uranyl acetate and lead citrate and examined on a Phillips BioTwin electron microscope (FEI, UK Ltd, Cambridge, UK). Electron microscope (EM) images of cross sections of fascicles within each nerve (areas ranging between 6732 and 47215 µm²) were analysed by eye by a trained, blinded observer using Image Tool 3.0 (UTHSCSA, USA). The total area of the nerve sections and the percentage of the nerve area occupied by nerve fascicles were calculated in order to investigate any differences between normal and laminitic digital nerves that might reflect oedema and therefore affect the quantification of axon density. The number and axon diameter of intact myelinated fibers was calculated as well as the percentage of damaged myelinated fibers, defined as those with a severe disruption of the myelin sheath and/or axonal degeneration. Myelin sheath thickness was measured and G-ratio of axons was calculated by dividing the axonal diameter by the total diameter of axon plus myelin sheath.

The proportion of A-fibers with continuous Schwann cell cytoplasm (an abnormal morphological feature previously described by Court et al., 2004) was also determined. C-fibers were identified as small-diameter unmyelinated fibers, surrounded by Schwann cell cytoplasm. The total number of C-fibers was calculated as well as the percentage of solitary unmyelinated fibers and the number of unmyelinated fibers per Remak bundle. All analysis was carried out on identity-concealed samples.

2.3. Immunohistochemistry

DRG from cervical segments 8 (forelimb innervation) and 4 (non-forelimb innervation) from the same horse were obtained post-mortem from the five laminitic horses and four control horses. The tissue was snap frozen and embedded in OCT embedding matrix (Cell Path plc. Powys, Wales, UK). Cryostat sections of C8 DRGs (15 µm) were thaw-mounted on poly-L-lysine slides (Merck-BDH).

DRG sections were pre-incubated for 1h at room temperature in 0.1 M PBS, pH 7.4, buffer containing 0.2% Triton X-100, 2% fish skin gelatin and 10% normal goat serum; and then incubated overnight at 4 °C with primary antibodies diluted in the same buffer. For co-localisation of the peptide NPY or ATF3 with the myelinated cell marker neurofilament 200 kDa (NF-200) (Lawson and Waddell, 1991; Michael et al., 1999), or either of the unmyelinated cell markers, peripherin or isolectin B4 (IB4) (Goldstein et al., 1991; Michael and Priestley, 1999), antisera/lectin were used at the following concentrations: rabbit anti-NPY (1:250; Peninsula Laboratories Inc, Belmont, CA, USA); rabbit anti-ATF3 (1:300; Santa Cruz Biotechnology, Santa Cruz, CA, USA); mouse monoclonal anti-NF-200 (1:400; clone N52; Sigma); mouse monoclonal anti-peripherin (1:250; Chemicon International, Harlow, UK); IB4 from *Bandeiraea simplicifolia* (1:400; Sigma). Sections were then washed in buffer and incubated at room temperature for 2 hours with Alexafluor 488-labeled goat anti-mouse IgG (1:500; Molecular Probes Europe BV, The Netherlands), Alexafluor 568-labeled goat anti-rabbit IgG (1:1000; Molecular Probes Europe BV, The Netherlands) or

Alexa Fluor 488-labeled streptavidin (1:200). Three washes in 0.1M PBS were performed before the addition of To-Pro3 cyanine nucleic acid stain (Molecular Probes Europe BV, The Netherlands). Three final washes in 0.1 M PBS were conducted before cover-slipping with Vecta-Shield (Vector Laboratories, Burlingame, CA, USA). Control sections were processed as above omitting the primary reagents.

Observations were made and sections photographed on an Olympus microscope equipped for epifluorescence. All counts of profiles labelled for immunopositive cells were performed by the same observer (who was blinded to sample treatment) on randomly selected, 15 µm sections of DRG from each of the animals in each group. Every sixth section was selected to ensure that measurements were taken only once for each cell. Results were expressed as the proportion of labelled profiles per total number of single or double-labelled profiles from all sections, 95% confidence intervals (CI) are indicated.

2.4. Western blots

C4 and C8 DRG were taken from laminitic horses (n=3). Whole lysate preparations were prepared by homogenising tissue in 20 volumes of Laemmli lysis buffer (Tris (tris-hydroxymethylaminoethane, 50 mM, pH 7.4), 5% mercaptoethanol and 2% sodium dodecyl sulphate (SDS)), boiled for 5 min and frozen. Western blotting was carried out as described previously (Garry et al., 2005). Blots were incubated with rabbit polyclonal primary antibodies to ATF3 (1:200 Santa Cruz Biotechnology, Santa Cruz, CA, USA) and detected by peroxidase-linked secondary antibody and enhanced chemiluminescence. The ubiquitous housekeeping enzyme glyceraldehyde-3-phosphate dehydrogenase (GAPDH, 1:750, Chemicon) was monitored as a control for protein level normalisation. Quantitative densitometry analysis of protein bands was performed using the ScanAnalysis (Elsevier) program.

2.5 Statistical analyses

Linear mixed-effect models were used to determine any differences between laminitic and controls horses in the frequency of lifting the forelimb and time spent at the back of the box, in order to account for the repeated sampling of the same horses (Pinheiro and Bates, 2000). The ID of the horse that the samples came from was entered as a random effect. Laminitic/control, time point in experiment and time of day were entered as fixed effects. Prior to analysis of the forelimb lifting results, the data were square root-transformed to achieve normalisation of the residuals.

For the analysis of percentages of damaged A-fibers, A-fibers with Schwann cell cytoplasm and solitary unmyelinated fibers, only one meaned value per horse was obtained, and therefore repeated sampling has not taken place. Repeated measures of mean axon diameters in myelinated and unmyelinated fibers and thickness of myelin sheath were taken in both control and laminitic horses. Therefore, linear-mixed effect models were also used to determine any differences between (i) mean axon diameters in myelinated and unmyelinated fibers; (ii) thickness of myelin sheath from laminitic and control horses. Multiple measurements per horse were also taken of the number of fibers per Remak bundle but as the data were integers, differences in the number of fibers per Remak bundle were analysed using generalised linear mixed-effect models with Poisson errors. Only a single measurement per horse of the percentage of damaged A fibers; A fibers with continuous Schwann cell cytoplasm and solitary unmyelinated fibers were taken, therefore simple logistic regressions were employed to determine the differences between control and laminitic horses.

Differences in total nerve area occupied by fascicles between normal and laminitic horses were investigated using a Student's t-test, and differences in the percentage of nerve area occupied by fascicles between normal and laminitic horses by general linear models with binomial errors. Any differences in the proportion of labelled profiles were assessed by χ^2 analysis. Mann-Whitney non-parametric tests were used to analyse fiber density. Immunoblot data were analysed using a matched pair t-test. All analyses were carried out in S-PLUS 6.0 (Insightful, Seattle, USA) and SigmaStat 2.03 (SPSS Inc., USA). In all cases $p < 0.05$ was

taken to indicate statistical significance, and degrees of freedom associated with any tests are denoted by subscripts.

3. Results

3.1. Radiographic abnormalities associated with laminitis were seen in all the laminitic horses.

The horses clinically diagnosed with laminitis displayed radiographic evidence of this disease when the radiographs were evaluated objectively (Butler et al., 2000). The angle between the dorsal hoof wall and the dorsal surface of the distal phalanx was increased when compared to normal values (Fig 1a). The mean (\pm SD) values from the laminitic group were $8.2^\circ \pm 3.0^\circ$ (normal values $-0.86^\circ \pm 2.4^\circ$) (Cripps and Eustace, 1999). Assessment of the D distance between the extensor process of the distal phalanx and the coronary band also showed a marked increase in the laminitic group ($D = 16.4\text{mm} \pm 4.9\text{mm}$) when compared to normal values ($4.1\text{mm} \pm 2.17\text{mm}$) (Cripps and Eustace, 1999). Histological sections of laminitic tissue also indicated inflammatory changes (Fig. 1d).

3.2. Laminitic horses display quantifiable abnormal behaviours

3.2.1. Data Analysis: Forelimb lifting

When considering overall data, laminitic horses show a statistically significant increase in the mean square root frequency of forelimb lifting ($F_{1,12}=11.5$, $p=0.005$; Fig. 2a) adjusted for time spent standing compared to control horses recorded in the same environment over the same time period. Fluctuations in the frequency of this behaviour occurred in both groups over the period of observation, but the pattern of such changes did not differ significantly between control and laminitic horses ($F_{1,140}=3.6$, $p=0.059$).

3.2.2. Data analysis proportion of time spent at the back of the box

Laminitic horses spent significantly more time at the back of the box than control horses ($F_{1,12}=6.1$, $p=0.03$). There was no difference between the 2 groups in how behaviour altered throughout the study ($F_{1,148}=0.2$, $p=0.683$) (Fig. 2b). Time spent at the back of the box was markedly higher in laminitics than in controls at both 06:00h and 22:00h but not at 14:00h (days 2 & 3), when assessing individual sample point data. This effect is not seen at 14:00h on day one, probably because drug administration regimes were not well-established at this time.

3.3. Distinct morphological abnormalities in both myelinated and unmyelinated peripheral nerve fibers innervating the hoof, in the lateral digital nerve of laminitic horses.

The lateral digital nerves at the level of the proximal sesamoid bone were examined from both normal and laminitic horses. A mean of 11.65% (range 8.75-14.35) of the total fascicle area from each nerve section was analysed. EM analysis of three randomly selected fascicles per lateral digital nerve revealed morphological differences in both the myelinated and unmyelinated fiber populations in laminitic compared to non-laminitic horses (Table 2). Abnormalities in the shape of surviving axons and disruption of the myelin sheath, with accumulation of lipid droplets and myelin debris were observed. The most obvious quantitative feature appeared to be a significant reduction in the number of both unmyelinated (-43.2%) and myelinated fibers (-34.6%) per unit area in laminitic compared to control horses ($p=0.016$). In order to eliminate the possibility that any nerve oedema could artefactually lead to the appearance of reduced fiber density, morphometric analyses were carried out to measure the percentage area of nerve sections occupied by fascicles and total nerve area in normal compared to laminitic horses. No significant differences in the mean percentages were identified ($t_4=-0.91$, $p=0.414$) between normals 37% (95% CI: 35.7-38.1) and laminitics 41% (39.5-42.0). No significant differences in total nerve area were identified between normal and laminitic horses ($t_4=-0.43$, $p=0.692$). Further abnormalities were a significant decrease in the

number of unmyelinated nerve fibers per Remak bundle ($F_{1,7}=20.7, p=0.003$) together with an increase in the percentage of solitary unmyelinated fibers in laminitics compared to normal horses ($\chi^2_1= 35.7, p<0.001$, Fig. 3; Table 2b). The percentage of morphologically damaged myelinated fibers was significantly higher in laminitic horses when compared to normal horses ($\chi^2_1= 31.5, p<0.001$, Fig. 3; Table 2 a). Finally, the proportion of myelinated fibers with continuous Schwann cell cytoplasm was significantly higher in the laminitic horses ($\chi^2=338.4, p<0.001$). No significant differences in myelin thickness or G-ratios were identified in laminitic compared to normal horses ($F_{1,7}<0.5, p>0.311$).

3.4. The neuronal injury marker ATF3 is selectively expressed in sensory neurons innervating the forelimb in laminitic horses.

Using immunohistochemical analysis of the DRG cell population, we assessed the presence of ATF3 in comparison with the expression of NF-200 and either IB4 or peripherin. ATF3 expression was significantly increased in NF-200-positive C8 DRG cells from laminitic horses (n=3), where 67% (15 sections, 304 cells, CI 58.8-69.9) of NF-200-positive DRG cells co-expressed ATF3, while only 10% (15 sections, 345 cells, CI 6.9-13.5) of NF-200-positive DRG cells in control horses (n=3) co-localised ATF3 ($\chi^2_1=208, p<0.001$), (Fig. 4). There was a significantly increased expression of ATF3 in IB4-positive C8 DRG cells in laminitic horses (n=3), where 54% (5 sections, 53 cells, CI 41.5-67.3) of IB4-positive C8 DRG cells co-localised ATF3 compared with 9% (5 sections 56 cells, CI 3.9-19.3) in control horses (n=3) ($\chi^2_1 = 24.5, p<0.001$). There was also a significantly increased proportion of peripherin-positive DRG cells that were positive for ATF3 in C8 DRG cells from laminitic horses, (n=3), where 57% (9 sections, 115 cells, CI 47.6-65.4) of peripherin-positive cells co-expressed ATF3 compared with 21% (9 sections, 127 cells CI 13.8-29.4) showing double-labelling in control horses (n=3) ($\chi^2_1 = 30.9, p=0.001$; Fig. 4).

Accordingly, immunoblot analysis revealed a significant increase ($p<0.05$) in ATF3 expression (expressed as mean percentage of GAPDH expression) in C8 DRG (38.9% (28.7-49.1) in comparison to low levels in the control C4 DRG (4.6% (-0.5-9.7) (Fig.4b, f). The numbers of cells expressing NF-200, IB4 or peripherin were unaltered in laminitic DRG compared to normal horses (696 compared to 575 NF-200-IR cells, $n=32$ sections, 115 compared to 127 peripherin-IR cells, $n=9$ sections, 56 compared to 53 IB4-IR cells, $n=5$ sections, in normal compared to laminitic horses, respectively).

3.5. Laminitis is associated with a distinctive pattern of expression of Neuropeptide Y (NPY) in sensory neurons.

A significant increase in NPY immunoreactivity (NPY-IR) was observed in the C8 DRG of laminitic horses, where 77% (17 sections, 271 cells, CI 72.0-82.3) of NF-200-positive cells showed NPY-IR co-localisation, compared to only 10% (17 sections, 351 cells CI 15.7-24.7) in control horses ($\chi^2_1 = 193$, $p<0.001$; Fig. 4).

4. Discussion

Damage to sensory nerves has been linked to abnormal pain and heightened sensitivity to touch in a variety of clinical and experimental studies. In this study, we have quantified for the first time abnormal behaviours associated with equine laminitis which are indicative of a hypersensitive sensory state. Additionally, we provide novel evidence for changes associated with nerve damage in the sensory nerves innervating the forelimb in laminitic horses, which are consistent with those reported in previously characterised neuropathic pain states.

4.1. The laminitic horses included in this study have digital pathology.

Assessment of the radiographs from the laminitic horses identified pedal bone displacement (rotation or distal displacement) associated with laminar tearing. It was not possible to perform radiographic assessment of the control horses due to ethical and health and safety

limitations, therefore data were compared to well established normal data (Cripps and Eustace, 1999). Chronic inflammatory changes were also observed (Fig. 1f, g) which have been previously shown to associate with sensory nerve losses in the skin (Lacomis et al., 1997; Tseng et al., 2006).

4.2. Laminitic horses display chronically altered behaviour.

We have quantified two behavioural changes associated with laminitis, which are suggestive of a chronic hypersensitive neuropathic pain state, characterised by the development of allodynia, hyperalgesia and spontaneous pain. Forelimb lifting represents an abnormal, *de novo* behaviour associated with laminitis, being at low levels or absent in the clinically normal horse. The overall scores for frequency of forelimb lifting were significantly greater in laminitic than control horses. Additionally, laminitic horses spent more time positioned towards the back of the box, a retiring behaviour that has also been associated with acute post-surgical limb pain (Price et al, 2003). This behaviour may represent a reluctance to engage in the external environment and preference to remain withdrawn from surroundings. The differences in retiring behaviour (percentage of time spent at the back of the box) showed the appearance of a marked cyclical pattern, although this was not seen with forelimb lifting, weight bearing when walking or general demeanour. The pattern apparent in time at the back of the box observations may correspond to NSAID dosing times, external environmental stimuli or possibly an intrinsic diurnal rhythm. This emphasises the need for behavioural testing at a number of regular intervals in order to correctly reveal specific changes. Moreover, the consistent deviations from normal behaviour over the three-day period confirmed that the NSAID analgesic regime was not consistently effective. When individual time points were evaluated, marked differences from control horses were consistently seen at 06:00h and 22:00h observations.

4.3. Abnormal hoof sensory nerve morphology in laminitic horses is consistent with that reported in damaged peripheral nerves in neuropathic pain states.

Two types of sensory receptor have been identified in the equine foot. Lamellated corpuscles, similar to Pacinian corpuscles, found primarily in the solar dermis of the heel, are low-threshold mechanoreceptors, which transmit their input via rapidly conducting, myelinated A-fibers (Bowker et al., 1993). Additionally, numerous naked nerve endings containing the neuropeptide, calcitonin gene related peptide (CGRP)-like immunoreactivity and other sensory neuropeptides such as substance P, neurokinin A and PHI (peptide histidine-isoleucine) were detected in the dermis of the dorsal hoof wall and sole (Bowker et al., 1995). Those containing CGRP are associated with nociception (Schmidt, 1981) and transmit via slowly conducting C-fibers. Axons from the hoof nociceptors and low-threshold mechanoreceptors as well as sympathetic fibres innervating the vasculature contribute to the sensory digital nerve.

Changes in any of these could potentially contribute to the etiology of the chronic laminitic pain state. To understand the mechanisms underlying laminitis pain and the incomplete response to anti-inflammatory analgesics, it is important to establish whether axonopathic changes may contribute. In laminitic horses, EM analysis identified marked decreases in myelinated and unmyelinated fiber numbers per unit area of digital nerve. This is unlikely to be due to nerve oedema as there were no significant differences between nerve section areas and fascicle areas in normal and laminitic horses. The marked increase in the number of solitary, unmyelinated fibers, may reflect demyelinated A-fibers, or an absence of guiding pathways for regenerating C-fibers (Bester et al., 1998). These morphological changes are consistent with those in laboratory neuropathic pain models, such as chronic constriction injury (Gautron et al., 1990, Basbaum et al., 1991, Gabay and Tal, 2004; Micu et al., 2006), crush injury (Lozeron et al., 2004), photochemically-induced ischemia (Yu et al., 2000) and diabetic neuropathy (Sima et al., 1988; Llewelyn et al., 1991; Elias et al., 1998; Kalichman et

al., 1998), thereby supporting our hypothesis that peripheral nerve damage may contribute to laminitis pain.

Functional changes in the injured peripheral nerve have also been described in neuropathic pain models. The loss of large fibers in nerves from laminitic horses is important as part of the behavioural changes in neuropathic pain states may result from the loss of spinal inhibitory controls exerted indirectly by these afferents (Basbaum et al., 1991). On the other hand, damage to both A and C-fibers appears to be necessary for the establishment of hyperalgesia and allodynia (Yu et al., 2000; Gabay and Tal, 2004). Electrophysiological studies further suggest that ectopic discharges in both spared C- and A-fibers may be important in maintaining neuropathic pain (Kajander and Bennett, 1992; Ali et al., 1999; Gabay and Tal, 2004).

4.4. Sensory neurons of the forelimb in laminitic horses show characteristic changes associated with peripheral nerve injury

Following peripheral nerve damage, phenotypic changes occur in primary sensory neurons that may contribute to mediating central sensitisation (Hokfelt et al., 1994; Tsujino et al., 2000). We assessed whether key neurochemical changes in sensory neurons of laminitic horses are similar to those in rodent neuropathic pain models. The numbers of DRG cells expressing anatomical markers NF-200, peripherin or IB4 were unaltered. Following nerve crush injury, peripherin increases transiently in large DRG cells (Wong and Oblinger, 1990). However, that model is associated with sensory loss (Bester et al., 1998) rather than the hypersensitivity seen here, as in other neuropathic and inflammatory pain states, where indeed peripherin expression is not upregulated (Facer et al., 2007, Renton et al., 2003, Rodriguez Parkitna et al., 2006).

Neuronal expression of ATF3, which is normally minimal, is upregulated after peripheral nerve injury and so acts as a marker of nerve injury (Tsujino et al., 2000). The clear expression of ATF3 in NF-200, peripherin or IB4-positive sensory neurons of laminitic horses, indicates neuronal damage to both A and C-fibers matching our observations of abnormal nerve morphology. These findings suggest that primary afferent injury associated with laminitis arises locally from the damage caused by hoof pathology, rather than from systemic disease, since ATF3 expression is low in neurons of unaffected limbs. Ischemia and ischemia/reperfusion are established causes of ATF-3 expression (Hai et al., 1999), so the ischemia-reperfusion injury of the digit thought to underlie acute laminitis (Hood et al., 1993) may also be involved in neuronal damage. We also found upregulated expression of NPY in large NF-200-positive DRG cells from laminitic horses, paralleling observations in other neuropathic pain models (Wakisaka et al. 1991, 1992; Noguchi et al., 1993; Kashiba et al., 1994; Nahin et al., 1994; Munglani et al., 1995; Rittenhouse et al., 1996; Ma and Bisby, 1998; Wallace et al., 2003).

The novel findings reported here suggest that pathological changes occurring during laminitis bring about a chronic pain state with a neuropathic component. Although the mechanisms underlying the pathogenesis of laminitis remain to be fully elucidated, it is apparent that the early stages of laminitis are associated with vasoconstriction of the digital microvasculature (Peroni et al., 2006) and inflammation (Belknap et al., 2007). Indeed, such pathological events can result in nerve damage (e.g. Yu et al., 2000; Zimmermann, 2001; Moalem and Tracey, 2006) and may thereby play a part in laminitis pain through the transition from acute inflammatory pain to a chronic syndrome with a neuropathic pain component.

Future studies will address the cellular and molecular mechanisms involved in the chronic laminitic pain state. These changes may be responsible, at least in part, for the limited efficacy of currently used anti-inflammatory therapy. The administration of anti-neuropathic agents may therefore achieve better pain management and improved quality of life in horses suffering from refractory laminitis.

Acknowledgements

This work was supported by the International League for the Protection of Horses (ILPH) (EJ and RAE) and The Royal (Dick) School of Veterinary Medicine. The BBSRC supported a summer studentship (ECS). We thank staff at Easter Bush Veterinary Centre for animal husbandry, Craig Penicuik for expert dissection, Gordon Goodall for histological preparation, Susan Kempson for histological assessment of laminitic hooves, Steven Mitchell for electron microscopy expertise, Linda Wilson, Biomedical Sciences and Shona Johnston, Centre for Inflammation Research, for confocal expertise, Colin Warwick for illustrations and Rod Else for helpful suggestions. This study would not have been possible without the support of the owners of the horses, for which we are grateful.

References

- Ali Z, Ringkamp M, Hartke TV, Chien HF, Flavahan NA, Campbell JN, Meyer RA. Uninjured C-fiber nociceptors develop spontaneous activity and alpha-adrenergic sensitivity following L6 spinal nerve ligation in monkey. *J Neurophysiol* 1999; 81:455-66.
- Basbaum AI, Gautron M, Jazat F, Mayes M, Guilbaud G. The spectrum of fiber loss in a model of neuropathic pain in the rat: an electron microscopic study. *Pain* 1991; 47: 359-67.
- Belknap JK, Giguere S, Pettigrew A, Cochran AM, Van Eps AW, Pollitt CC. Lamellar pro-inflammatory cytokine expression patterns in laminitis at the developmental stage and at the onset of lameness: innate vs. adaptive immune response. *Equine Vet J* 2007; 39: 42-7.
- Bester H, Allchorne AJ, Woolf J. Recovery of C fiber-Induced Extravasation Following Peripheral Injury in the Rat. *Exp Neurol* 1998; 154: 628-36.
- Bowker RM, Brewer AM, Vex KB, Guida LA, Linder KE, Sonea IM, Stinson AW. Sensory receptors in the equine foot. *Am J Vet Res* 1993; 54: 1840-4.
- Bowker RM, Linder K, Sonea IM, Guida LA. Sensory nerve fibers and receptors in equine distal forelimbs and their potential roles in locomotion. *Equine Vet J Suppl* 1995; 18: 141-6.
- Butler J, Colles C, Dyson S, Kold, S, Poulos P. Foot, Pastern and Fetlock. In: *Clinical Radiology of the Horse*. Oxford: Blackwell Science, 2000. pp. 27-60.

486 Court FA, Sherman DL, Pratt T, Garry EM, Ribchester RR, Cottrell DF, Fleetwood-Walker
487 SM, Brophy PJ. Restricted growth of Schwann cells lacking Cajal bands slows conduction in
488 myelinated nerves. *Nature* 2004; 431: 191-5.

489 Cripps P.J and Eustace R.A. Radiological measurements with reference to laminitis. *Equine*
490 *Vet Journal* 1999; 31: 427-32

491 Cummins TR, Black JA, Dib-Hajj SD, Waxman SG. Glial-derived neurotrophic factor
492 upregulates expression of functional SNS and NaN sodium channels and their currents in
493 axotomized dorsal root ganglion neurons. *J Neurosci* 2000; 20: 8754-61.

494 Dyson SJ. The diagnosis of laminitis. In: Dyson S and Hall M, editors. *Diagnosis and*
495 *management of lameness in the horse*. Missouri: WB Saunders, 2003. pp. 329.

496 Facer P, Casula MA, Smith GD, Benham CD, Chessell IP, Bountra C, Sinisi M, Birch R,
497 Anand P. Differential expression of the capsaicin receptor TRPV1 and related novel receptors
498 TRPV3, TRPV4 and TRPM8 in normal human tissues and changes in traumatic and diabetic
499 neuropathy. *BMC Neurol* 2007; 7: 11-22.

500 Gabay E, Tal M. Pain behaviour and nerve electrophysiology in the CCI model of neuropathic
501 pain. *Pain* 2004; 110: 354-60.

502 Garry EM, Delaney A, Blackburn-Munro G, Dickinson T, Moss A, Nakalembe I, Robertson
503 DC, Rosie R, Robberecht P, Mitchell R, Fleetwood-Walker SM. Activation of p38 and p42/44
504 MAP kinase in neuropathic pain: Involvement of VPAC(2) and NK(2) receptors and
505 mediation by spinal glia. *Mol Cell Neurosci* 2005; 30: 523-37.

506 Gautron M, Jarat F, Ratinahirana H, Hauw JJ, Guilbaud G. Alterations in myelinated fibers in
507 the sciatic nerve of rats after constriction: possible relationships between the presence of
508 abnormal small myelinated fibers and pain-related behaviour. *Neurosci Lett* 1990; 111: 28-33.

509 Gerring EL, Lees P, Taylor JB. Pharmacokinetics of phenylbutazone and its metabolites in the
510 horse. *Equine Vet J* 1981; 13: 152-7.

511 Goldstein ME, House SB, Gainer H. NF-L and peripherin immunoreactivities define distinct
512 classes of rat sensory ganglion cells. *J Neurosci Res* 1991; 30: 92-104.

513 Hai T, Wolfgang CD, Marsee DK, Allen AE, Sivaprasad U. ATF3 and stress responses. *Gene*
 514 *Expr* 1999; 7: 321-35.

515 Herthel D, Hood DM. Clinical presentation, diagnosis, and prognosis of chronic laminitis.
 516 *Vet Clin North Am Equine Pract* 1999; 15: 375-94.

517 Hokfelt T, Zhang X, Wiesenfeld-Hallin Z. Messenger plasticity in primary sensory neurons
 518 following axotomy and its functional implications. *Trends Neurosci* 1994; 17: 22-30.

519 Hood DM, Grosenbaugh DA, Mostafa MB, Morgan SJ, Thomas BC. The role of vascular
 520 mechanisms in the development of acute equine laminitis. *J Vet Intern Med* 1993; 7: 228-34.

521 Hood DM. The pathophysiology of developmental and acute laminitis. *Vet Clin North Am*
 522 *Equine Pract* 1999; 15: 321-43.

523 Kajander KC, Bennett, GJ. Onset of a painful peripheral neuropathy in rat: A partial and
 524 differential deafferentation and spontaneous discharge in A β and A δ primary afferent
 525 neurons. *J Neurophysiol* 1992; 68: 734-44.

526 Kashiba H, Noguchi K, Ueda Y, Senba E. Neuropeptide Y and galanin are coexpressed in rat
 527 large type A sensory neurons after peripheral transection. *Peptides* 1994; 15: 411-6.

528 Lawson SN, Waddell PJ. Soma neurofilament immunoreactivity is related to cell size and
 529 fiber conduction velocity in rat primary sensory neurons. *J Physiol* 1991; 435: 41-63.

530 Lacomis D, Giuliani MJ, Steen V, Powell HC. Small fiber neuropathy and vasculitis. *Arthritis*
 531 *Rheum* 1997; 40: 1173-7.

532 Ma W, Bisby MA. Partial and complete sciatic nerve injuries induce similar increases of
 533 neuropeptide Y and vasoactive intestinal peptide immunoreactivities in primary sensory
 534 neurons and their central projections. *Neuroscience* 1998; 86: 1217-34.

535 Michael GJ, Averill S, Shortland PJ, Yan Q, Priestley JV. Axotomy results in major changes
 536 in BDNF expression by dorsal root ganglion cells: BDNF expression in large trkB and trkC
 537 cells, in pericellular baskets, and in projections to deep dorsal horn and dorsal column nuclei.
 538 *Eur J Neurosci* 1999;11: 3539-51.

539 Michael GJ, Priestley JV. Differential expression of the mRNA for the vanilloid receptor
540 subtype 1 in cells of the adult rat dorsal root and nodose ganglia and its downregulation by
541 axotomy. *J Neurosci* 1999;19:1844-54.

542 Micu I, Jiang Q, Coderre E, Ridsdale A, Zhang L, Woulfe J, Yin X, Trapp BD, McRory JE,
543 Rehak R, Zamponi GW, Wang W, Stys PK. NMDA receptors mediate calcium accumulation
544 in myelin during chemical ischaemia. *Nature* 2006; 439: 988-92.

545 Moalem G and Tracey DJ. Immune and inflammatory mechanisms in neuropathic pain. *Brain*
546 *Res Rev* 2006; 51: 240-64.

547 Munglani R, Bond A, Smith GD, Harrison SM, Elliot PJ, Birch PJ, Hunt SP. Changes in
548 neuronal markers in a mononeuropathic rat model relationship between neuropeptide Y, pre-
549 emptive drug treatment and long-term mechanical hyperalgesia. *Pain* 1995; 63: 21-31.

550 Nahin RL, Ren K, De Leon M, Ruda M. Primary sensory neurons exhibit altered gene
551 expression in a rat model of neuropathic pain. *Pain* 1994; 58: 95-108.

552 Noguchi K, De Leon M, Nahin RL, Senba E, Ruda MA. Quantification of axotomy-induced
553 alteration of neuropeptide mRNAs in dorsal root ganglion neurons with special reference to
554 neuropeptide Y mRNA and the effects of neonatal capsaicin treatment. *J Neurosci Res* 1993;
555 35: 54-66.

556 Peroni JF, Moore JN, Noschka E, Grafton ME, Aceves-Avila M, Lewis SJ, Robertson TP.
557 Predisposition for venoconstriction in the equine laminar dermis: implications for equine
558 laminitis. *J Appl Physiol* 2006; 100: 759-763.

559 Pinheiro JC, Bates D M. Mixed-effects models in S and S-plus. New York: Springer-Verlag,
560 2000.

561 Pollitt CC, Pass MA, Pollitt S. Batimastat (BB-94) inhibits matrix metalloproteinases of
562 equine laminitis. *Equine Vet J Suppl* 1998; 26: 119-24.

563 Pollitt CC. Medical management of laminitis. In: Dyson S and Hall M, editors. *Diagnosis and*
564 *management of lameness in the horse*. Missouri: WB Saunders, 2003. pp. 329-31.

565 Price, J, Catriona, S, Welsh, EM, Waran, NK. Preliminary evaluation of a behaviour-based
566 system for assessment of post-operative pain in horses following arthroscopic surgery. *Vet*
567 *Anaesth Analg* 2003; 30(3): 124-137.

568 Reitmann TR, Stauffacher M, Bernasconi P, Auer JA, Weishaupt MA. The association
569 between heart rate, heart rate variability, endocrine and behavioural pain measures in horses
570 suffering from Laminitis. *J Vet Med* 2004; 51: 218-25.

571 Renton T, Yiangou Y, Baecker PA, Ford AP, Anand P. Capsaicin receptor VR1 and ATP
572 purinoreceptor P2X3 in painful and nonpainful human tooth pulp. *J Orofac Pain* 2003; 17:
573 245-50.

574 Rittenhouse PA, Marchand JE, Chen J, Kream RM, Leeman SE. Streptozotocin-induced
575 diabetes is associated with altered expression of peptide-encoding mRNAs in rat sensory
576 neurons. *Peptides* 1996; 17: 1017-22.

577 Rodriguez Parkitna J, Korostynski M, Kaminska-Chowanec D, Obara I, Mika J, Przewlocka
578 B, Przewlocki R. Comparison of gene expression profiles in neuropathic and inflammatory
579 pain. *J Physiol Pharmacol* 2006; 57: 401-14.

580 Schmidt RG. Somatovisceral sensibility. In: Schmidt RG, editor. *Fundamentals of sensory*
581 *Physiology*. New York: Springer-Verlag, 1981. pp. 81-125.

582 Stashak TS. Diagnosis of Lameness. In: Stashak TS, editor. *Adam's Lameness in Horses*. 5th
583 ed. Philadelphia: Lea and Febiger, 2002. pp. 649-652

584 Swanson TD. Clinical presentation, diagnosis, and prognosis of acute laminitis. *Vet Clin*
585 *North Am Equine Pract* 1999; 15: 311-9.

586 Tseng M-T, Hsieh S-C, Shun C-T, Lee K-L, Pan C-L, Lin W-M, Lin Y-H, Yu C-L, Hsieh S-
587 T. Skin denervation and cutaneous vasculitis in systemic lupus erythematosus. *Brain* 2006;
588 129: 977-85.

589 Tsujino H, Kondo E, Fukuoka T, Dai Y, Tokunaga A, Miki K, Yonenobu K, Ochi T, Noguchi
590 K. Activating transcription factor 3 (ATF3) induction by axotomy in sensory and
591 motoneurons: A novel neuronal marker of nerve injury. *Mol Cell Neurosci* 2000; 15: 170-82.

Wakisaka S, Kajander KC, Bennett GJ. Increased neuropeptide Y (NPY)-like immunoreactivity in rat sensory neurons following peripheral axotomy. *Neurosci Lett* 1991; 124: 200-3.

Wakisaka S, Kajander KC, Bennett GJ. Effects of peripheral nerve injuries and tissue inflammation on the levels of neuropeptide Y-like immunoreactivity in rat primary afferent neurons. *Brain Res* 1992; 598: 349-52.

Wallace CJ, Cottrell DF, Brophy PJ, Fleetwood-Walker SM. Focal lysolecithin-induced demyelination of peripheral afferents results in neuropathic pain behavior that is attenuated by cannabinoids. *J Neurosci* 2003; 23: 3221-33.

Wong J, Oblinger MM. Differential regulation of peripherin and neurofilament gene expression in regenerating rat DRG neurons. *J Neurosci Res* 1990; 27: 332-41.

Woolf CJ, Salter MW. Neuronal plasticity: Increasing the gain in pain. *Science* 2000; 288: 1765-8.

Xiao HS, Huang QH, Zhang FX, Bao L, Lu YJ, Guo C, Yang L, Huang WJ, Fu G, Xu SH, Cheng XP, Yan Q, Zhu ZD, Zhang X, Chen Z, Han ZG, Zhang X. Identification of gene expression profile of dorsal root ganglion in the rat peripheral axotomy model of neuropathic pain. *Proc Natl Acad Sci U S A* 2002; 99: 8360-5.

Yu W, Kauppila T, Hultenby K, Persson JK, Xu XJ, Weisenfeld-Hallin Z. Photochemically-induced ischemic injury of the rat sciatic nerve: a light- and electron microscope study. *J Peripher Nerv Syst* 2000; 5: 209-17.

Zimmermann M. Pathobiology of neuropathic pain. *Eur J of Pharmacol* 2001; 429: 23-37.

Fig. 1. (a) Latero-medial radiograph of laminitic equine digit showing rotation and vertical displacement (“sinking”) of the third phalanx relative to normal anatomy. Lines represent the standardised methods for measuring displacement (D) which is the distance (mm) between the proximal limit of the dorsal hoof wall and the extensor process of the distal phalanx, and the rotation angle (α) which is the angle between the dorsal surface of the distal phalanx and the dorsal surface of hoof wall (Cripps and Eustace, 1999). (b) Haemotoxylin/Eosin (H&E) stained histological section of the intact lamellar distal phalangeal apparatus in a normal horse showing the normal appearance of dermal (arrowhead) and epidermal (arrow) laminae x10. (c) H&E stained histological section showing disruption and separation of the secondary epidermal (arrow) and secondary dermal (arrowhead) laminae in acute laminitis x10. (d) Chronic laminitis. Mild inflammation in the laminar dermis, with small foci of lymphocytes in a perivascular location (arrows). H&E, original magnification x20. (e) Chronic laminitis. Blood accumulation in the inner stratum medium. H&E, original magnification x 4. Scale bars (b-e) = 100 μ m.

Fig. 2. Quantifiable pain behaviours in laminitic horses compared to control horses. Behavioural indices were recorded in laminitic horses (▲, solid line) (n=7) and clinically normal (□, dashed line) horses (n=7) over a period of 3 days, with 1-hour observations at 06:00 hrs, 14:00 hrs and 22:00 hrs. Phenylbutazone was administered each day at 08:00h and 20:00h with supplementary acepromazine at 08:00h, 16:00h and 24:00h. (a) Forelimb lifting frequency adjusted for total time standing, expressed as lifts/min. When considering overall data, laminitic horses show a statistically significant increase in the mean square root frequency of forelimb lifting (\pm SE) adjusted for time spent standing compared to control horses recorded in the same environment over the same time period. (b) Proportion of time spent at the back of the box (away from the entrance), expressed as a percentage of time (\pm SE). Laminitic horses show a marked increase in the overall proportion of time spent at the back of the box, with marked differences from control horses at 06:00h and 22:00h.

Fig. 3. Reduced myelinated and unmyelinated fiber density associated with laminitis. (a) Electron microscopy images of digital nerve from a normal horse. Arrows indicate intact, normal myelinated fibers. Arrowheads indicate clustered unmyelinated fibers in Remak bundles. (b) Electron microscopy images of digital nerve from a laminitic horse displaying reduced myelinated fiber density (arrows), lower numbers of C-fibers per Remak bundle, as well as increased numbers of solitary fibers (arrowheads) and increased collagen-filled space, compared to normal horse. Scale bars, 5 μ m.

Fig. 4. (a-e) Immunohistochemical co-localisation of DRG neuronal subtype markers (NF-200 and peripherin, green) with neuronal injury marker, ATF3 or neuropeptide Y (NPY) (red) in C8 DRG (which receives forelimb innervation) of laminitic (a,d,e), or control horses (b) and co-localisation of NF-200 (green) with ATF3 (red) in C4 DRG (not associated with forelimb innervation) from the same horse (c). (a) In laminitic horses, there was an increased expression of ATF3 (red) in NF-200-positive DRG cells (green) compared to C8 DRG control (non-laminitic) horse (b) and C4 DRG from laminitic horse (c). Laminitic horses show expression of ATF-3 (red) in peripherin-positive (green) in DRG cells (d), while control horses do not (data not shown). Additionally, there was increased co-localisation of NPY (red) and NF-200 (green) in C8 DRG cells of laminitic horses (e) compared to control horses, where there was normally only sparse NPY expression (data not shown). Scale bars, 100 μ m. White arrows show co-localised immunopositive cells. Open arrows show cell marker (NF-200 or peripherin)-positive cells lacking co-localisation. (f) Typical immunoblots of whole DRG lysates of laminitic horses (n=3), show clear ATF3 expression in C8 but not C4 DRG. Levels of the housekeeping enzyme, GAPDH (lower blots) were unchanged.

Table 1. Details of horses used in the study. Sex abbreviations: MN –male, neutered; M- male, intact, F- female. Treatment abbreviations: PBZ – phenylbutazone; F – flunixin; A – aspirin, ACP – acepromazine; SS – solar supports; RF – remedial farriery; FT – foot trimming; NG – nitroglycerin (vasodilatory therapy); R – rehydration therapy; T – Trilostane (modifier of steroidogenesis); P – procaine penicillin + neomycin sulphate.

Estimated weight range for laminitic horses: 250-550 kg; age range: 6-21 years. Control horses used were 3 females, 3 neutered male, and an intact male. Estimated weight range for control horses: 350-600 kg; age range: 8-19 years.

Horse Group/No.	Sex	Time from onset	Possible precipitating/concurrent conditions	Prior treatments
Laminitic 1	MN	1 month	Obese, increased liver enzymes	PBZ, ACP, NG, SS, RF, A, T
Laminitic 2	F	2 months	Obese	PBZ, ACP, RF, FT, SS
Laminitic 3	F	Recurrent >1 year	Obese	PBZ
Laminitic 4	F	2 days	None known	ACP, SS, F, A
Laminitic 5	MN	Recurrent >1 year	Grain overload	PBZ, F, SS, R, P
Laminitic 6	M	Recurrent >1 year	Obese	PBZ, RF, SS, NG, FT
Laminitic 7	MN	Recurrent >1 year	Access to rich pasture	PBZ, SS
Laminitic 8	MN	1 month	None known: prior history unknown	PBZ, FT, ACP, NG, SS, RF, A
Laminitic 9	MN	Recurrent >1 year	Euthanasia requested for chronic condition	PBZ, FT, RF
Laminitic 10	MN	Recurrent >1 year	None known: prior history unknown	PBZ, FT
Laminitic 11	MN	Recurrent >1 year	None known: prior history unknown	PBZ, RF
Laminitic 12	MN	Recurrent >1 year	None known: prior history unknown	PBZ, RF

Table 2. Summary of myelinated and unmyelinated nerve fiber characteristics in normal and laminitic horses. Statistical significance is indicated by asterisks (* P value of <0.05, ** P<0.01, *** P<0.001; Linear mixed effects models, Mann Whitney test – Mean no. of fibers per 100 μm^2 -). Values are expressed as mean \pm SEM.


a) Myelinated fibers

Nerve fiber characteristics	Normal Horses n=4	Laminitic Horses n=5
Mean no. of fibers per 100 μm^2	0.52 (\pm 0.04)	0.34 (\pm 0.02) *
Mean percentage of damaged A fibers	16.40 (\pm 2.75)	30.08 (\pm 5.67) ***
Mean percentage of A fibers with continuous (>40%) Schwann cell cytoplasm	17.45 (\pm 1.63)	72.46 (\pm 5.85) ***
Mean axon diameter (μm)	5.38 (\pm 0.1)	5.08 (\pm 0.1)
Mean thickness of myelin sheath (μm)	1.06 (\pm 0.03)	1.09 (\pm 0.04)

b) Unmyelinated fibers

Nerve fiber characteristics	Normal Horses n=4	Laminitic Horses n=5
Mean no. of fibers per 100 μm^2	5.77 (\pm 0.53)	3.28 (\pm 0.31) *
Mean no. of fibers per Remak bundle	2.75 (\pm 0.07)	2.09 (\pm 0.03) **
Mean percentage of solitary unmyelinated fibers	30.14 (\pm 2.33)	38.06 (\pm 5.15) ***
Mean axon diameter (μm)	1.36 (\pm 0.01)	1.28 (\pm 0.01)

835 Fig. 1.
836
837



838 **Fig. 2.**
839
840






Fig. 3.



892 Fig. 4.
893



894